# Development of a new wound assessment form

Wound assessment is a routine component of caring for patients with any type of wound. To date, there is little agreement about how assessment is carried out and recorded and several published audits have identified that in many instances it is done inconsistently. A project group met to develop and agree a new wound assessment tool which, it is suggested, may form the basis for agreeing a minimum dataset. The layout of the form is specifically designed to facilitate ease of use in combination with digital pen technology, making it quick and simple to both input and audit data.

## Jacqueline Fletcher

## **KEY WORDS**

Wound assessment Audit National benchmark Best Practice

ounds are a major source of morbidity to patients and a major cost to hospitals and community healthcare providers (Posnett et al, 2009). As the UK population ages, the number of patients with both acute and chronic wounds increases, with costs to the NHS estimated to be in the range of £2.3-£3.1 billion (at 2005/2006) for chronic wounds alone (Posnett and Franks, 2007). In addition, the complications associated with wounds place an additional burden on resources. Surgical site infections (SSIs) (which account for up to 20% of hospital-acquired infections [HAIs], National Institute for Health and

Clinical Excellence [NICE], 2008) can have a significant impact on resources. Surgical site infections may range from spontaneously limited wound discharge within 7–10 days of an operation, to a life-threatening postoperative complication. It is reported that over one-third of postoperative deaths are related to SSIs (NICE, 2008). Other clinical outcomes of SSIs include poor scars that are cosmetically unacceptable, such as those that are spreading, hypertrophic or keloid, persistent pain and itching, restriction of movement, particularly when over joints, and

a significant impact on emotional wellbeing (NICE, 2008). Surgical site infections are estimated to incur additional costs of between £814 and £6,626 per patient (NICE, 2008), and at least double the length of hospital stay (Health Protection Agency [HPA], 2009), depending on the type of surgery and severity of the infection.

Every patient with a wound has a right to expect a good minimum standard of care, regardless of the cause of their wound or where that care is delivered. When a patient with a wound is managed inappropriately,

#### BOX I

# Members of the project team

- Mark Collier, Lead Nurse, Tissue Viability, United Lincoln Hospitals NHS Trust – Acute
- >> Dr Caroline Dowsett, Nurse Consultant, Tissue Viability, Newham Primary Care Trust
- >>> Jacqueline Fletcher, Senior Professional Tutor, Department of Wound Healing, Cardiff University and Principal Lecturer, Tissue Viability, University of Hertfordshire
- >> Brenda King, Nurse Consultant, Tissue Viability, Sheffield Community
- >> Kathryn Vowden, Nurse Consultant, Tissue Viability, Bradford Teaching Hospitals NHS Foundation Trust and University of Bradford
- >> Trudie Young, Lecturer in Tissue Viability, Bangor University

Jacqueline Fletcher is Professional Tutor, Department of Dermatology and Wound Healing, Cardiff and Principal Lecturer, Tissue Viability, University of Hertfordshire they can suffer from failure to heal, resulting in the wound being present longer than is necessary and an increased risk of complications. Posnett and Franks (2008) stated that a high proportion of chronic wounds remain unhealed for long periods and for almost certainly longer than necessary. Such ineffective management can result not only in prolonged patient suffering, but also increased costs to healthcare organisations.

In the Best Practice Statement Optimising Wound Care (Harding et al, 2008), the authors suggest that in order to provide a good standard of care, a structured approach is required to assessment, diagnosis and management of patients with wounds, and that assessment is fundamental to planning care. The Best Practice Statement maintains that, 'A thorough patient assessment should be carried out by a skilled and competent practitioner adhering to local and national guidelines, when appropriate, at all levels in the service'. However, assessment (and recording of the assessment) is an area of practice which is often carried out poorly or sporadically (Dowsett, 2009). Dowsett (2009) in a study of community nurses' knowledge and practice, identified that at baseline only 42% of patients had a wound assessment form completed, which is consistent with audit findings elsewhere (Ashton and Price, 2006; McIntosh and Ousey, 2008).

Although most clinicians would suggest that they do perform an assessment, this is frequently not evident from their documentation. Previously, an audit of 83 sets of leg ulcer documentation identified that the use of a specific assessment chart significantly increased the likelihood of appropriate data being collected and recorded, and where no chart was used, information was difficult to find and often omitted (Fletcher. 2001). Although in almost half of the notes audited there was inadequate information, it would appear from the auditors' reports that the patients were generally receiving appropriate care, however, this is a subjective view

## BOX 2

## List of descriptors included in the forms

- Name (These would be the basic details on an addressograph label)
- >> Age/DOB
- >> GP/consultant
- >> Address/ward/department
- >> Date of assessment
- >> Signature of assessor

## Type of wound

#### History of wound

#### Location of wounds

May give actual locations or include a body map/diagram

#### Measurements

#### Tissue description

Usually with some indication of the % attributed to each type

## Symptom description

Pain

Exudate

Odour

#### Surrounding skin

#### Specific risk assessments,

E.g. ankle brachial pressure index (ABPI), pressure ulcer risk and grade

## Factors that may delay healing

#### Referrals

May also include a section on treatments: these may be coded especially if there is a local formulary — so a limited number of products — or may be open space for free text

Objective of care

Cleansing solution

Primary dressing

Secondary dressing

**Padding** 

Bandage/tape

Frequency of dressing change

Re-assessment date

and would be difficult to substantiate at a later point in time. In Lord Darzi's report, the *Next Stage Review* (Department of Health [DH], 2008), he firmly sets quality at the heart of the NHS, stating that we need to be clear about what high quality care looks like, and that in order to improve we need to be able to measure and understand exactly what we do. Furthermore, this lack of recording

information in a systematic way makes it difficult to maintain continuity of care, particularly in the community setting where many different practitioners may be involved in the care of the same patient (Dowsett, 2009).

Despite recommendations for formalised wound assessment (Harding et al, 2008), there are no

## BOX 3

# Symptom descriptions

Pain could use a range of pain rating systems

- Intensity
- >> Numerical rating scale, e.g 0-5 or 1-10
- >> Visual rating scale, e.g. smiling faces
- >> Verbal rating scale, e.g. none, mild, moderate severe

#### **Nature**

A range of descriptor words, e.g. sharp, stabbing, dull or a blank space to record the patient's description

#### Frequency

- Constant
- Procedural
- Incident
- Intermittent

#### Exudate

Volume; a variety of descriptors:

- >> None, scant, moderate, high or none, low, moderate high very high
- Dry/none, slight (weekly dressing change), moderate (2/3 weekly dressing change), copious (daily or more changes) +, ++, +++
- May also ask: is the level increasing/decreasing/static

## Colour

- >> Serous, serous sanguinous, sanguinous, pus
- >> Clear, blood-stained, pus
- >> Yellow, green, red, cream
- >> Clear/amber, cloudy/milky or creamy, pink or red, green, yellow or brown, grey or blue (European Wound Management Association [EWMA], 2007)

#### Viscosity

>> Thick, stringy, thin/runny

#### Odour

- +,++,++
- >> Baker and Haig descriptors, e.g. not evident at arm's length, similar to TELER

recommendations for what should be included within such a form. Furthermore, there is no agreement on how to describe the indicators used within the forms. Given the current drivers to improve quality of care, measure standardised outcomes and the message from Darzi to 'get the basics right every time', which must include fundamentals such as wound assessment, this is immensely problematic (DH, 2008).

## Development of the standardised form

As part of a project to develop a standardised wound assessment form (Box I notes the members of the project team) for use with digital pen technology (Vowden, 2009), a review was carried out of 33 assessment forms (17 generic and 16 leg ulcer forms). It was apparent that most areas collected similar data but the way this was done varied considerably. Some forms appeared to strive for

simplicity and to be on one side of A4, collecting only minimal information, for example, the descriptor for pain would simply say 'yes/no', while others collected much more comprehensive data with, for example, information on the intensity, nature, frequency and duration of pain.

The list of descriptors included in the forms can be seen in *Box 2*, with an example of how the descriptors may be expanded within the various forms in *Box 3*.

The assessment charts were reviewed by the author who compiled a spreadsheet of common terms and the frequency with which they occurred. Following this review, the project group met to attempt to determine key factors which should be included in a wound assessment. form. There was considerable discussion around every individual factor and reference was made throughout to key documents, such as the World Union of Wound Healing Societies (WUWHS) document on wound exudate (WUWHS, 2007) (Box 4 lists the other documents that were referred to). Where possible, existing descriptors were used, although in some instances these provoked considerable discussion. An example of this would be the descriptors proposed for wound moisture levels, i.e. dry, moist, wet, saturated and leaking. While these have clearly been agreed by the WUWHS expert panel, there was considerable debate about the first three (dry, moist, wet) relating to the wound bed condition, and the last two, saturated and leaking, appearing to relate to an assessment of the dressing condition. Although this may seem to be purely semantics, the project group were keen to ensure that there was minimal possibility for misinterpretation or misunderstanding.

Once the key criteria and appropriate descriptors had been agreed by the project group, consideration was given to the presentation of the information. As the form is primarily designed to be used

## BOX 4

## **Key documents consulted**

- >> EWMA (2006) Position Document. Management of wound infection
- >> EWMA (2005) Position Document. Identifying criteria for wound infection
- >> EWMA (2002) Position Document. Pain at wound dressing changes
- WUWHS (2008) Principles of best practice. Wound infection in clinical practice: An international consensus
- WUWHS (2007) Principles of best practice. Wound exudate and the role of dressings
- WUWHS (2004) Principles of best practice. Minimising pain at dressing related procedures. A consensus document

with digital pen technology, tick boxes were preferable as they are quicker for staff completing the form. The ordering of the information had to be logical, for example, descriptors indicating progress or deterioration.

The first two pages of the assessment form are composed of primarily demographic data which would only be captured on one occasion, and a body map — a feature which all of the project team felt should be included as it enables a quick visual location of the wound and easy numerical identification if more than one wound is present. The following three pages cover standard wound assessment data (i.e. details

# **National Wound Assessment Form**

Wound Details			
Date of assessment	Date of assessment	Date of assessment	Date of assessment
DDMMYYYY	DDMMYYYY	D D M M Y Y Y Y	DDMMYYYY
Wound Number	Wound Number	Wound Number	Wound Number
1 2 3 4 5	1 2 3 4 5	1 2 3 4 5	1 2 3 4 5
6 7 8 9 10	6 7 8 9 10	6 7 8 9 10	6 7 8 9 10
Has wound been traced?	Has wound been traced?	Has wound been traced?	Has wound been traced?
Yes No	Yes No	Yes No	Yes No
If yes: Max length mm	If yes: Max length mm	If yes: Max length mm	If yes: Max length mm
Max width mm	Max width mm	Max width mm	Max width mm
Type of wound	Type of wound	Type of wound	Type of wound
Pressure ulcer 1 2 3 4	Pressure ulcer 1 2 3 4	Pressure ulcer 1 2 3 4	Pressure ulcer 1 2 3 4
Leg ulcer	Leg ulcer	Leg ulcer	Leg ulcer
Fungating lesion	Fungating lesion	Fungating lesion	Fungating lesion
Diabetic foot ulcer	Diabetic foot ulcer	Diabetic foot ulcer	Diabetic foot ulcer
Surgical wound	Surgical wound	Surgical wound	Surgical wound
Traumatic wound	Traumatic wound	Traumatic wound	Traumatic wound
Burn / Scald	Burn / Scald	Burn / Scald	Bum / Scald
Cellulitis (Weeping)	Cellulitis (Weeping)	Cellulitis (Weeping)	Cellulitis (Weeping)
Other (specify below)	Other (specify below)	Other (specify below)	Other (specify below)
Other/Comments	Other/Comments	Other/Comments	Other/Comments

Figure 1. National Wound Assessent form.

Wound Details			
Tissue Type (% of each type of tissue in wound)	Tissue Type (% of each type of tissue in wound)	Tissue Type (% of each type of tissue in wound)	Tissue Type (% of each type of tissue in wound)
Necrotic (black/brown)	Necrotic (black/brown)	Necrotic (black/brown)	Necrotic (black/brown)
Sloughy (cream/yellow)	Sloughy (cream/yellow)	Sloughy (cream/yellow)	Sloughy (cream/yellow)
Granulating (red)	Granulating (red)	Granulating (red)	Granulating (red)
Epithelialising (pink)	Epithelialising (pink)	Epithelialising (pink)	Epithelialising (pink)
Infection clinical signs Present Yes No If yes, is it: Local Systemic Both Indicators of local infection	Infection clinical signs Present Yes No If yes, is it: Local Systemic Both Indicators of local infection	Infection clinical signs Present Yes No If yes, is it: Local Systemic Both Indicators of local infection	Infection clinical signs Present Yes No If yes, is it: Local Systemic Both Indicators of local infection
Cellulitis  Abscess/pus	Cellulitis  Abscess/pus	Cellulitis  Abscess/pus	Cellulitis  Abscess/pus
Increased pain	Increased pain	Increased pain	Increased pain
Increased exudate	Increased exudate	Increased exudate	Increased exudate
Malodour	Malodour	Malodour	Malodour
Delayed healing / deterioration	Delayed healing / deterioration	Delayed healing / deterioration	Delayed healing / deterioration
Friable granulation tissue/bleeds easily			
Pocketing at wound base			
Date swab sent	Date swab sent	Date swab sent	Date swab sent
DDMMYYY	Y D D M M Y Y Y Y	DDMMYYYY	DDMMYYYY
Swab results	Swab results	Swab results	Swab results
MRSA	MRSA	MRSA	MRSA
Pseudomonas	Pseudomonas	Pseudomonas	Pseudomonas
Haemolytic strep A	Haemolytic strep A	Haemolytic strep A	Haemolytic strep A
Anaerobes	Anaerobes	Anaerobes	Anaerobes
Staph. aureus	Staph. aureus	Staph. aureus	Staph. aureus
Wound moisture levels	Wound moisture levels	Wound moisture levels	Wound moisture levels
Dry	Dry	Dry	Dry
Moist	Moist	Moist	Moist
Wet	Wet	Wet	Wet
Saturated	Saturated	Saturated	Saturated
Leaking	Leaking	Leaking	Leaking

Figure 2. National Wound Assessent form.

Pain frequency Continuous	Pain frequency Continuous	Pain frequency Continuous	Pain frequency Continuous
Intermittent	Intermittent	Intermittent	Intermittent
At dressing change	At dressing change	At dressing change	At dressing change
Post dressing	Post dressing	Post dressing	Post dressing
None	None	None	None
Does patient report wound odour as a problem?	Does patient report wound odour as a problem?	Does patient report wound odour as a problem?	Does patient report wound odour as a problem?
Yes No	Yes No	Yes No	Yes No
Is the wound reported to be:			
Deteriorating	Deteriorating	Deteriorating	Deteriorating
Static	Static	Static	Static
Improving	Improving	Improving	Improving
Healed	Healed	Healed	Healed
Treatment objectives	Treatment objectives	Treatment objectives	Treatment objectives
Tick all that apply			
Protecting granulation / epithelialisation			
Debridement of necrosis or slough			
Manage bacterial burden	Manage bacterial burden	Manage bacterial burden	Manage bacterial burden
Re-hydration of wound bed			
Managing exudate	Managing exudate	Managing exudate	Managing exudate
Protection of surrounding skin			
Managing wound pain	Managing wound pain	Managing wound pain	Managing wound pain
Minimise odour	Minimise odour	Minimise odour	Minimise odour

Figure 3. National Wound Assessent form.

that occurred most commonly in the review) and allow for four assessments (Figures 1, 2 and 3). Categories include:

- >> Date of assessment
- Wound number (if more than one wound present)
- Has the wound been traced?
- >> Type of wound
- Duration of the wound
- >> Tissue type and percentage
- >> Clinical signs of infection
- >> Indicators of infection
- >> Swab sent and result
- ▶ Wound moisture levels
- >> Surrounding skin condition
- >> Wound pain (level and frequency)
- ▶ Wound odour
- Current status of the wound (deteriorating, static, improving, healed)
- >> Treatment objectives.

The following two pages of the form identify treatment details, such as dressing used, cleansing carried out, additional fixation, with the final page allowing for any additional notes to be made.

It is acknowledged by the project group that there is little research to support inclusion of any of the criteria identified within the form as reliable indicators of wound progress, other than the measurement of the wound which can be used to determine probability of healing (Cardinal et al, 2008). It must also be noted that even this has been questioned, as both the technique and accuracy of the various methods of measuring wound area and volume differ considerably (Jessop, 2005; Langemo et al, 2008; Little et

al, 2009). However, it appears from both the project group's experience and the review of forms in current use, that practitioners rely on the indicators used (i.e. tissue type, size, etc) to provide information to help set objectives and measure progress. This broad experiential-based development process relates closely to the views of Leaper (2009), which challenge the tyranny of the randomised controlled trial (RCT) within wound care.

This is the first draft of the form but the group believe that it is the first time that consensus has been reached (albeit by a small group) on both the content and layout of a wound assessment form, thus giving it peer validation. The initial form has been piloted within a clinical area (Vowden,

2009) and some minor amendments have been made to layout and wording (e.g. ordering of information and size of boxes).

While the project group do not propose that the form should become a 'national standard' without further consultation, it is a positive step to see agreement across both acute and community settings on the minimum dataset that is required. It is hoped that the form will focus discussion and be a initial step towards developing a national benchmark, which, as the Best Practice document *Optimising Wound Care* recommends, should be in place and auditable so that every patient has a minimum standard of wound care.

Copies of the form are available online from: <a href="https://www.e-fficient.co.uk">www.e-fficient.co.uk</a>

#### **Acknowledgements**

The project group and development of the form were kindly sponsored by Coloplast UK. Technical development of the form was by Longhand data.

## References

Ashton J, Price P (2006) Survey comparing clinicians' wound healing knowledge and practice. *Br J Nurs Tissue Viability Supplement* **15(19)**: S18–S26

Cardinal M, Eisenbud DE, Philips T, Harding K (2008) Early healing rates and wound area measurements are reliable predictors of later complete wound closure. *Wound Rep Regen* 16(1):19–22

Department of Health (2009) Transforming community services: ambition, action, achievement. DH, London. Available online at: <a href="www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/">www.dh.gov.uk/en/PublicationsAndStatistics/Publications/PublicationsPolicyAndGuidance/</a>
DH 101425

Dowsett C (2009) Use of TIME to improve community nurses' wound care knowledge and practice. *Wounds UK* 5(3): 14–21

European Wound Management Association (EWMA) (2002) Position Document: *Pain at wound dressing changes*. London: MEP Ltd. Available online at: <a href="http://ewma.org/fileadmin/user\_upload/EWMA/pdf/Position\_Documents/2002/Spring\_2002\_English\_.pdf">http://ewma.org/fileadmin/user\_upload/EWMA/pdf/Position\_Documents/2002/Spring\_2002\_English\_.pdf</a>

European Wound Management Association (EWMA) (2005) Position Document:

Identifying criteria for wound infection.
London: MEP Ltd. Available online at: <a href="http://ewma.org/fileadmin/user\_upload/EWMA/">http://ewma.org/fileadmin/user\_upload/EWMA/</a>
<a href="pdf/Position\_Documents/2005\_Wound\_Infection\_/English\_pos\_doc\_final.pdf">http://ewma.org/fileadmin/user\_upload/EWMA/</a>
<a href="pdf/Position\_Documents/2005\_Wound\_Infection\_Jenglish\_pos\_doc\_final.pdf">http://ewma.org/fileadmin/user\_upload/EWMA/</a>
<a href="pdf/Position\_Documents/2005\_Wound\_Infection\_Jenglish\_pos\_doc\_final.pdf">http://ewma.org/fileadmin/user\_upload/EWMA/</a>
<a href="pdf/Position\_Documents/2005\_Wound\_Infection\_Jenglish\_pos\_doc\_final.pdf">http://ewma.org/fileadmin/user\_upload

European Wound Management Association (EWMA) (2006) Position Document:

Management of wound infection. London: MEP
Ltd. Available online at: http://ewma.org/fileadmin/user\_upload/EWMA/pdf/Position\_Documents/2006/English\_pos\_doc\_2006.pdf

Fletcher J (2001) An audit of documentation to evaluate the implementation of leg ulcer guidelines across Hertfordshire wounds.

10th European Conference on Advances in Wound Management, Dublin

Harding KG, et al (2008) Best Practice Statement Optimising Wound Care. Wounds UK, Aberdeen. Available online at: <u>www.</u> <u>wounds-uk.com/downloads/BPS\_Optimising.pdf</u>

Health Protection Agency (HPA) (2009) Healthcare Associated Infections in England: 2008–2009 Report. HPA, London. Available online at: <a href="www.hpa.org.uk/web/HPAweb&HPAweb&HPAweb&HPAwebStandard/HPAweb\_C/1252326221795">www.hpa.org.uk/web/HPAweb&HPAweb&HPAweb\_C/1252326221795</a>

Jessop RL (2005) What is the best method for assessing the rate of wound healing? A comparison of 3 mathematical formulas. *Adv Skin Wound Care* 19(138): 140–6

Langemo D, Aderson J, Hanson D, Hunter S, Thompson P (2008) Measuring wound length, width and area: which technique? *Adv Skin Wound Care* 21(1): 42–5

Leaper D (2009) Evidence-based wound care in the UK. *Int Wound Journal* **6(2)**: 89–91

Little C, McDonald J, Jenkins MG, and McCarron P (2009) An overview of techniques used to measure wound area and volume. J Wound Care 18(6): 250–3

McIntosh C, Ousey K (2008) A survey of nurses' and podiatrists' attitudes, skills and knowledge of lower extremity wound care. *Wounds UK* 4(1): 59–68

National Institute for Health and Clinical Excellence (2008) Surgical Site Infection. Prevention and treatment of surgical site infection. NICE Clinical Guideline 74. NICE, London. Available online at: <a href="http://guidance.nice.org.uk/CG74/Guidance/pdf/English">http://guidance.nice.org.uk/CG74/Guidance/pdf/English</a>

Posnett J, Franks PJ (2007) The cost of skin breakdown and ulceration in the UK. In: The Smith and Nephew Foundation (2007) *Skin Breakdown the Silent Epidemic*. Smith and Nephew Foundation. Hull

Posnett J, Franks PJ (2008) The burden of chronic wounds in the UK. *Nurs Times* **104(3)** Chronic Wound Supplement

## **Key points**

- Wound assessment is a routine component of caring for patients with any type of wound.
- Every patient with a wound has a right to expect a good minimum standard of care.
- To develop a standardised wound assessment form for use with digital pen technology (Vowden, 2009), a review was carried out of 33 assessment forms.
- This is the first time that consensus has been reached (albeit by a small group) on both the content and layout of a wound assessment form, thus giving it peer validation

Posnett J, Gottrup F, Lundgren H, Saal G (2009) The resource impact of wounds on health care providers in Europe. *J Wound Care* 18(4): 154–61

Vowden K (2009) A new methodology for data collection. Wounds UK Conference, Harrogate

World Union of Wound Healing Societies (WUWHS) (2008) Principles of best practice: Wound infection in clinical practice. An international consensus. London: MEP Ltd. Available online at: <a href="https://www.mepltd.co.uk/pdf/Wound%20Inf%20">www.mepltd.co.uk/pdf/Wound%20Inf%20</a> S&N\_English\_WEB.pdf

World Union of Wound Healing Societies (2007) Principles of best practice. Wound exudate and the role of dressings. London: MEP Ltd. Available online at: <a href="https://www.mepltd.co.uk/download\_pdfs/consensus%20">www.mepltd.co.uk/download\_pdfs/consensus%20</a> exudate%20Eng%2007.pdf

World Union of Wound Healing Societies (2004) Principles of best practice: Minimising pain at dressing related procedures. A consensus document. London: MEP Ltd. Available online at: <a href="https://www.mepltd.co.uk/download\_pdfs/consensus%20pain%20Eng%2004.pdf">www.mepltd.co.uk/download\_pdfs/consensus%20pain%20Eng%2004.pdf</a>